

## Quantitive and qualitive peculiarities of the zooplankton and its formation in the East Georgia

Zelimkhan Kereselidze

[zelimkhan.kereselidze@tsu.ge](mailto:zelimkhan.kereselidze@tsu.ge)

Department of Biology, Faculty of Exact and Natural Science, Iv.Javakishvili Tbilisi State University,  
University str.2, 0143 Tbilisi, Georgia

Key words: Zooplankton, Eutrophication, Dominant forms, qualitative structure

Eutrophication process reveals in qualitative and qualitative structure of Zooplankton in reservoirs – are changed the total amount of species and the amount of dominant forms are changed, also varies the number and biomass of Zooplankton.

Last summer we took Planktonic material of Kumisi, Sion, Bazaleti and Tabatskuri lakes. We processes them by the Camerally way and compared with other water materials of the same time. The following picture had been revealed:

In Kumisi reservoir the amounts of Zooplankton species have been decreased threefold in 50 years, also dominant forms have been changed. Currently reigning evribiont Copepods, the amount and biomass of Zooplankton are reduced.

In Sioni reservoir we see different situation: dominant forms aren't changed so far and quantitative structure had been varied in a small frameworks.

In Bazaleti lake has been emerged Eutrophication sings: Dominant forms amount had been decreased and also the amount and biomass are changed. The same situation is in Tabatskuri lake.

### Reference:

1. В. И. Жадин. Жизнь пресных вод СССР. Том 3. М.,Л.. 1950 г.
2. А. В. Монаков, Труды биологической станции „Борок“, выпуск 3, 1958 г.
3. Е. Ф. Мануйлова, Труды биологической станции „Борок“, выпуск 3, 1958 г.
4. Н. А. Дзюбан, Труды шестого совещания по проблемам биологии внутренних вод, М..., Л... 1959 г.
5. Л. И. Лебедева, Бюллетень МОИП отдел биологический №6, 1961 г.
6. Г.Б. Мельников, Гидробиологический журнал, том 2. №2. 1966 г.